

02 07 - 00

PTO/SB/05 (4/98)

Please type a plus sign (+) inside this box → ☐Approved for use through 09/30/2000. OMB 0651-0032  
Patent and Trademark Office: U.S. DEPARTMENT OF COMMERCE

Under the Paperwork Reduction Act of 1995, no persons are required to respond to a collection of information unless it displays a valid OMB control number.

**UTILITY  
PATENT APPLICATION  
TRANSMITTAL**

(Only for new nonprovisional applications under 37 C.F.R. § 1.53(b))

Attorney Docket No. **ARC2914R1**First Inventor or Application Identifier **Stephen A. Berry**Title **See 1 in Addendum**Express Mail Label No. **EL523936456US****APPLICATION ELEMENTS**

See MPEP chapter 600 concerning utility patent application contents.

ADDRESS TO:

Assistant Commissioner for Patents  
Box Patent Application  
Washington, DC 20231

1. ☒ \* Fee Transmittal Form (e.g., PTO/SB/17)  
(Submit an original and a duplicate for fee processing)
2. ☒ Specification [Total Pages **33**]  
(preferred arrangement set forth below)
- Descriptive title of the Invention
  - Cross References to Related Applications
  - Statement Regarding Fed sponsored R & D
  - Reference to Microfiche Appendix
  - Background of the Invention
  - Brief Summary of the Invention
  - Brief Description of the Drawings (if filed)
  - Detailed Description
  - Claim(s)
  - Abstract of the Disclosure
3. ☒ Drawing(s) (35 U.S.C. 113) [Total Sheets **8**]
4. Oath or Declaration [Total Pages **4**]
- a. ☒ Newly executed (original or copy)
  - b. ☐ Copy from a prior application (37 C.F.R. § 1.63(d))  
(for continuation/divisional with Box 16 completed)
    - i. ☐ **DELETION OF INVENTOR(S)**  
Signed statement attached deleting  
inventor(s) named in the prior application,  
see 37 C.F.R. §§ 1.63(d)(2) and 1.33(b).

5. ☐ Microfiche Computer Program (Appendix)
6. Nucleotide and/or Amino Acid Sequence Submission  
(if applicable, all necessary)
- a. ☐ Computer Readable Copy
  - b. ☐ Paper Copy (identical to computer copy)
  - c. ☐ Statement verifying identity of above copies

**ACCOMPANYING APPLICATION PARTS**

7. ☒ Assignment Papers (cover sheet & document(s))
8. ☒ 37 C.F.R. § 3.73(b) Statement (when there is an assignee) ☒ Power of Attorney
9. ☐ English Translation Document (if applicable)
10. ☒ Information Disclosure Statement (IDS)/PTO-1449 ☒ Copies of IDS Citations
11. ☐ Preliminary Amendment
12. ☒ Return Receipt Postcard (MPEP 503)  
(Should be specifically itemized)
13. ☐ \* Small Entity Statement filed in prior application  
(PTO/SB/09-12) ☐ Status still proper and desired
14. ☐ Certified Copy of Priority Document(s)  
(if foreign priority is claimed)
15. ☐ Other: \_\_\_\_\_

\* NOTE FOR ITEMS 1 & 13 IN ORDER TO BE ENTITLED TO PAY SMALL ENTITY  
FEES, A SMALL ENTITY STATEMENT IS REQUIRED (37 C.F.R. § 1.27), EXCEPT  
IF ONE FILED IN A PRIOR APPLICATION IS RELIED UPON (37 C.F.R. § 1.28).

**16 If a CONTINUING APPLICATION, check appropriate box, and supply the requisite information below and in a preliminary amendment:**☐ Continuation ☐ Divisional ☐ Continuation-in-part (CIP) of prior application No. \_\_\_\_\_ / \_\_\_\_\_

Prior application information: Examiner \_\_\_\_\_

Group / Art Unit: \_\_\_\_\_

**For CONTINUATION or DIVISIONAL APPS only:** The entire disclosure of the prior application, from which an oath or declaration is supplied under Box 4b, is considered a part of the disclosure of the accompanying continuation or divisional application and is hereby incorporated by reference. The incorporation can only be relied upon for information inadvertently omitted from the submitted application parts.

**17. CORRESPONDENCE ADDRESS**☐ Customer Number or Bar Code Label**22921**or ☐ Correspondence address below

(Insert Customer Number or Bar code label here)

Name					
Address					
City	State	Zip Code			
Country	Telephone	Fax			

Name (Print/Type)	<b>Pauline Clarke</b>	Registration No. (Attorney/Agent)	<b>29,783</b>
Signature	<i>Pauline Clarke</i>	Date	<b>Feb 3, 2000</b>

Burden Hour Statement: This form is estimated to take 0.2 hours to complete. Time will vary depending upon the needs of the individual case. Any comments on the amount of time you are required to complete this form should be sent to the Chief Information Officer, Patent and Trademark Office, Washington, DC 20231. DO NOT SEND FEES OR COMPLETED FORMS TO THIS ADDRESS. SEND TO Assistant Commissioner for Patents, Box Patent Application, Washington, DC 20231.

**PATENT APPLICATION TRANSMITTAL LETTER**  
(Large Entity)

Docket No.  
ARC2914R1

TO THE ASSISTANT COMMISSIONER FOR PATENTS

Transmitted herewith for filing under 35 U.S.C. 111 and 37 C.F.R. 1.53 is the patent application of:

**Berry et al**

For: **STABLE NON-AQUEOUS SINGLE PHASE VISCOUS VEHICLES AND FORMULATIONS UTILIZING SUCH VEHICLES**

Enclosed are:

- ☒ Certificate of Mailing with Express Mail Mailing Label No. EL523936456US
- ☒ 8 sheets of drawings.
- ☐ A certified copy of a application.
- ☒ Declaration ☒ Signed. ☐ Unsigned.
- ☒ Power of Attorney
- ☒ Information Disclosure Statement
- ☐ Preliminary Amendment
- ☐ Other:

**CLAIMS AS FILED**

For	#Filed	#Allowed	#Extra	Rate	Fee
Total Claims	44	- 20 =	24	x \$18.00	\$432.00
Indep. Claims	4	- 3 =	1	x \$78.00	\$78.00
Multiple Dependent Claims (check if applicable) <input checked="" type="checkbox"/>					\$780.00
BASIC FEE					\$690.00
TOTAL FILING FEE					\$1,980.00

- ☐ A check in the amount of to cover the filing fee is enclosed.
- ☒ The Commissioner is hereby authorized to charge and credit Deposit Account No. 01-1173 as described below. A duplicate copy of this sheet is enclosed.
  - ☒ Charge the amount of \$1,980.00 as filing fee.
  - ☒ Credit any overpayment.
  - ☒ Charge any additional filing fees required under 37 C.F.R. 1.16 and 1.17.
  - ☐ Charge the issue fee set in 37 C.F.R. 1.18 at the mailing of the Notice of Allowance, pursuant to 37 C.F.R. 1.311(b).

Dated: Feb 3, 2000

  
Signature

CC:

Attachment to PTO/SB/05 (4/98) Utility Patent Application  
Transmittal

1. Stable Non-Aqueous Single Phase Viscous Vehicles and Formulations Utilizing Such Vehicles

STABLE NON-AQUEOUS SINGLE PHASE VISCOUS VEHICLES  
AND FORMULATIONS UTILIZING SUCH VEHICLES

5    Field of the Invention

          This invention relates to stable non-aqueous single phase  
biocompatible viscous vehicles capable of suspending beneficial agents and  
uniformly dispensing said agents at low flow rates and more particularly to  
stable uniformly mixed formulations of beneficial agents in stable non-  
10    aqueous single phase biocompatible viscous vehicles.

References

          The following references are referred to by numbers in brackets ([ ]) at  
the relevant portion of the specification.

- 15           1.     Wang, et al., *J. Parenteral Sci. Tech*, **42**: S4-S26 (1988).  
             2.     Desai, et al., *J. Am. Chem. Soc.*, **116**: 9420-9422 (1994).  
             3.     Chang, et al., *Pharm. Tech.*, 80-84 (Jan. 1996).  
             4.     Manning, et al., *Pharm. Res.*, **6**: 903-918 (1989).  
             5.     Hageman, *Drug Dev. Ind. Pharm*, **14**: 2047-2070 (1988).  
20           6.     Bell, et al., *Biopolymers*, **35**: 201-209 (1995).  
             7.     Zhang, et al., *Pharm. Res.* **12**: 1447-1452 (1995).  
             8.     PCT published application 98/00158  
             9.     PCT published application 98/16250  
             10.    Knepp, et al., *Pharm. Res.* **15** (7) 1090-1095 (1998).  
25           11.    PCT published application 98/00157  
             12.    PCT published application 98/00152  
             13.    U.S. Patent 5,540,912  
             14.    U.S. Patent 5,571,525  
             15.    U.S. Patent 5,512,293  
30           16.    PCT published application 96/40049  
             17.    Yu, et al., *J. Pharm. Sci.*, 85: 396-401 (1996).  
             18.    Mitchell, U.S. Patent No. 5,411,951 (1995).  
             19.    Brooks, et al., U.S. Patent No. 5,352,662 (1994)

20. Geller, L., U.S. Patent No. 3,869,549 (1975).
21. Larsen, et al., PCT Publication No. WO95/34285 (1995).
22. Knepp, et al., *J. Pharm. Sci. Tech*, **50**: 163-171 (1996).
23. U.S. Patent 5,614,221
- 5 24. U.S. Patent 4,594,108
25. U.S. Patent 5,300,302
26. U.S. Patent 4,588,614
27. U.S. Patent 4,310,516
28. U.S. Patent 5,635,213
- 10 29. EP 379,147

### Background of the Invention

Peptides, polypeptides, proteins and other proteinaceous substances (e.g., viruses, antibodies) collectively referred to herein as proteins, have great utility as pharmaceuticals in the prevention, treatment and diagnosis of disease. Proteins are naturally active in aqueous environments, thus the preferred formulations of proteins have been in aqueous solutions. However, proteins are only marginally stable in aqueous solutions. Thus, protein pharmaceuticals often have short shelf-lives under ambient conditions or require refrigeration. Further, many proteins have only limited solubility in aqueous solutions. Even when they are soluble at high concentrations, they are prone to aggregation and precipitation.

Because proteins can easily degrade, the standard method for delivering such compounds has been daily injections. Proteins can degrade via a number of mechanisms, including deamidations of asparagine and glutamine; oxidation of methionine and, to a lesser degree, tryptophan, tyrosine and histidine; hydrolysis of peptide bonds; disulfide interchange; and racemization of chiral amino acid residues [1-7]. Water is a reactant in nearly all of these degradation pathways. Further, water acts as a plasticizer, which facilitates unfolding and irreversible aggregation of proteins. Since water is a participant in almost all protein degradation pathways, reduction of aqueous

protein solution to a dry powder provides an alternative formulation methodology to enhance the stability of protein pharmaceuticals.

One approach to stabilizing proteins is to dry them using various techniques, including freeze-drying, spray-drying, lyophilization, and  
5 desiccation. Dried proteins are stored as dry powders until their use is required.

A serious drawback to drying of proteins is that often one would like to use proteins in some sort of flowable form. Parenteral injection and the use of drug delivery devices for sustained delivery of drug are two examples of the  
10 applications where one would like to use proteins in a flowable form. For injection, dried proteins must be reconstituted, adding additional steps which are time-consuming and where contamination may occur, and exposing the protein to potentially destabilizing conditions [7]. For drug delivery devices the protein formulations must be stable for extended periods of time at body  
15 temperature and maintain their flowability for the expected life of the device.

Solution formulations of proteins/peptides in non-aqueous polar aprotic solvents such as DMSO and DMF have been shown to be stable at elevated temperatures for long periods of time [8]. However, such solvent based formulations will not be useable for all proteins since many proteins have low  
20 solubility in these solvents. The lower the solubility of the protein in the formulation, the more solvent would have to be used for delivery of a specific amount of protein. Low concentration solutions may be useful for injections, but may not be useful for long term delivery at low flow rates.

Proteins have been formulated for delivery using perfluorodecalin [9,  
25 10], methoxyflurane [9], high concentrations in water [11], polyethylene glycol [12], PLGA [13, 14], ethylenevinylacetate/polyvinylpyrrolidone mixtures [15], PEG400/povidone [16]. However, these formulations were not shown to retain a uniform suspension of protein in viscous vehicle over long periods of time.

30 Many biologically active compounds degrade over time in aqueous solution. Carriers in which proteins do not dissolve but rather are suspended, can often offer improved chemical stability. Furthermore, it can be beneficial

to suspend the beneficial agent in a carrier when the agent exhibits low solubility in the desired vehicle. However, suspensions can have poor physical stability due to settling and agglomeration of the suspended beneficial agent. The problems with non-aqueous carriers tend to be exacerbated as the concentration of the active compound is increased.

Dispersing powdered proteins or peptides in lipid vehicles to yield parenteral sustained release formulations has been investigated [17-21]. The vehicles used were either various vegetable (sesame, soy, peanut, etc.) or synthetic oils (e.g., Miglyol) gelled with aluminum fatty acid esters such as aluminum stearates (mono-, di- or tri-), or with a polyglycerol ester. Although theoretically these vehicles might preclude solution denaturation and protect the drug from aqueous chemical degradation, the vehicles themselves are unstable at higher temperatures. The storage of liquid vegetable oils at body temperatures results in the formation of reactive species such as free fatty acids and peroxides (a process which is accelerated by the presence of traces of various metal ions such as copper or iron which can leach from some implantable devices). These peroxides not only adversely affect protein stability [22] but would be toxic when delivered directly to, for example, the central nervous system of a human or animal.

The sustained delivery of drugs has many advantages. Use of implantable devices assures patient compliance, since the delivery device is tamper-proof. With one insertion of a device, rather than daily injections, there is reduced site irritation, fewer occupational hazards for practitioners improved cost effectiveness through decreased costs of equipment for repeated injections, reduced hazards of waste disposal, and enhanced efficacy through controlled release as compared with depot injection. The use of implantable devices for sustained delivery of a wide variety of drugs or other beneficial agents is well known in the art. Typical devices are described, for example, in U.S. Patents Nos. 5,034,229; 5,057,318; 5,110,596; and 5,782,396. The disclosure of each of these patents is incorporated herein by reference.

For drug delivering implants, dosing durations of up to one year are not unusual. Beneficial agents which have low therapeutic delivery rates are prime candidates for use in implants. When the device is implanted or stored, settling of the beneficial agent in a liquid formulation can occur. This  
5 heterogeneity can adversely affect the concentration of the beneficial agent dispensed. Compounding this problem is the size of the implanted beneficial agent reservoir. Implant reservoirs are generally on the order of 25-250  $\mu$ l, but can be up to 25 ml.

Viscous formulations have been prepared using two separate  
10 components to be mixed with drug at use [23], thickening agents added to aqueous compositions [24], gelling agents added to aqueous drug solutions [25], porous textile sheet material [26], thickening agents with oleaginous material [27], viscous aqueous carrier for limited solubility drug [28], and extrudable elastic gels [29]. However, these formulations are mixed at use,  
15 contain aqueous components, use sheet matrices, or are delivered topically, orally, or intraduodenally.

Stability of formulations can be enhanced by freeze-drying, lyophilizing or spray-drying the active ingredient. The process of drying the active ingredient includes further advantages such as compounds which are  
20 relatively unstable in aqueous solution can be processed and filled into dosage containers, dried without elevated temperatures, and then stored in the dry state in which there are relatively few stability problems.

Pharmaceutical formulations, particularly parenteral products, should be sterilized after being sealed in the final container and within as short a time  
25 as possible after the filling and sealing have been completed. (See, for example Remington, Pharmaceutical Sciences, 15<sup>th</sup> ed. (1975)). Examples of sterilization techniques include thermal or dry-heat, aseptic, and ionized radiation. Combinations of these sterilization procedures may also be used to produce a sterile product.

30 There is a need to be able to deliver protein compositions to the body which are stable at body temperatures over extended periods of time to enable long term delivery of the protein. There is a need to be able to deliver



concentrations of proteins that are efficacious. There is a need for a novel non-aqueous formulation capable of homogeneously suspending proteins and dispensing such agents at body temperatures and low flow rates over extended periods of time.

5

### Summary of the Invention

The present invention provides stable single phase non-aqueous biocompatible viscous vehicles capable of forming uniform suspensions with proteins. The components of the viscous vehicle comprise at least two of  
10 polymer, surfactant, and solvent. The ratios of the components will vary depending on the molecular weight of the components and the desired viscosity of the final vehicle. Presently preferred component ratios are: polymer, about 5% to about 60%; solvent, about 30% to about 50%; and surfactant, about 5% to about 20%.

15 The present invention also provides stable formulations in which beneficial agents are uniformly suspended in stable single phase non-aqueous biocompatible viscous vehicles. In particular, the beneficial agents are formulated in the viscous vehicles at concentrations of at least about 0.1%, depending upon the potency of the beneficial agent. These stable  
20 formulations may be stored at the temperature appropriate for the beneficial agent, ranging from cold, to body temperature (about 37°C) for long periods of time (1 month to 1 year or more). In a preferred embodiment the formulation comprises about 0.1 to 50% (w/w) of beneficial agent, depending on the potency of the beneficial agent, the duration of treatment, and the rate of  
25 release for the drug delivery system.

These formulations are especially useful in implantable delivery devices for long term delivery (e.g., 1 to 12 months or longer) of beneficial agent at body temperature, preferably about 37°C. Thus, the present invention also provides for the delivery of said proteins to the body over  
30 extended period of time to enable long term delivery of the protein at low flow rates of about 0.3 to 100 µl/day, preferably about 0.3 to 4 µl/day for about a 6

month delivery period and preferably 5 to 8  $\mu$ l/day for about a 3 month delivery period.

In another aspect, the invention provides methods for preparing stable non-aqueous biocompatible formulations of a beneficial agent in a single phase viscous vehicle. Preferred formulations comprise about 0.1 to 50% (w/w) beneficial agent depending on the potency of the beneficial agent, the duration of treatment, and the rate of release from the delivery system.

In yet a further aspect, the invention provides methods for treating a subject suffering from a condition which may be alleviated by administration of a beneficial agent, said methods comprising administering to said subject an effective amount of a stable non-aqueous formulation comprising at least one beneficial agent uniformly suspended in a single phase viscous vehicle.

A further aspect of the invention is that non-aqueous single phase viscous vehicles containing beneficial agents are chemically and physically stable over a broad temperature range for long periods of time. The beneficial agents in the viscous vehicles are also chemically and physically stable over a broad temperature range for long periods of time. Thus, these formulations are advantageous in that they may be shipped and stored at temperatures below, at, or above room temperature for long period of time. They are also suitable for use in implantable delivery devices in which the formulation must be stable at body temperature for extended periods of time.

The formulations of the present invention also remain stable when delivered from implantable drug delivery systems. The beneficial agents have been shown to exhibit zero order release rates when delivered from implantable drug delivery systems at very low flow rates over extended periods of time.

#### Brief Description of the Drawings

Figure 1 shows the stability of hGH formulations of the present invention as determined at 37°C by reverse phase HPLC.

Figure 2 shows the stability of hGH formulations of the present invention as determined at 37°C by size exclusion chromatography.

Figure 3 shows the average release rate ( $\mu\text{l/day}$ ) of 10% (w/w) spray-dried lysozyme in formulations of the present invention.

Figure 4 shows the average release rate ( $\mu\text{l/day}$ ) of 10% (w/w) spray-dried hGH in a glycerol monolaurate/lauryl lactate/polyvinylpyrrolidone vehicle.

Figure 5 shows the average release rate ( $\mu\text{g/day}$ ) of 10% lysozyme in a lauryl alcohol/polyvinylpyrrolidone vehicle.

Figure 6 shows the average release rate ( $\mu\text{g/day}$ ) of 25% lysozyme in a glycerol monolaurate/lauryl lactate/polyvinylpyrrolidone vehicle.

Figure 7 shows the average release rate ( $\mu\text{g/day}$ ) of 33% lysozyme in a glycerol monolaurate/lauryl lactate/polyvinylpyrrolidone vehicle.

Figure 8 shows the average release rate ( $\mu\text{g/day}$ ) of 45% lysozyme in a glycerol monolaurate/lauryl lactate/polyvinylpyrrolidone vehicle.

#### Detailed Description of the Invention

The present invention is drawn to the unexpected discovery that uniformly suspending beneficial agents in non-aqueous single phase biocompatible viscous vehicles results in stable formulations which can be delivered at body temperature over an extended period of time at low flow rates. Previously known formulations of beneficial agents which are buffered aqueous or non-aqueous solutions which may or may not contain excipients do not provide formulations which can be uniformly dispensed at body temperatures at low flow rates over an extended period of time without exhibiting unacceptable amounts of aggregation or degradation of the formulation. The presently claimed formulations stabilize beneficial agents and can be stored at the temperature appropriate for the beneficial agent. The temperatures can range from cold (not exceeding  $8^{\circ}\text{C}$ ) to body temperature (about  $37^{\circ}\text{C}$ ) for long periods of time. These formulations are especially useful in implantable delivery devices for long term delivery (e.g., 1 to 12 months or longer) of drug at low flow rates and at body temperature, preferably about  $37^{\circ}\text{C}$ .

Standard beneficial agent formulations consist of dilute aqueous or non-aqueous solutions or suspensions. Drug stability is usually achieved by varying one or more of the following: pH, buffer type, ionic strength, excipients (EDTA, ascorbic acid, etc.) For these formulations, degradation pathways requiring water (hydrolysis, deamidation, racemization) cannot be fully stabilized. In the present invention, beneficial agents formulated in non-aqueous biocompatible single phase viscous vehicles containing for example, polyvinylpyrrolidone, vinyl acetate, and/or polyoxyethylenepolyoxypropylene block copolymers were shown to be chemically and physically stable. The viscosity of the formulation will depend upon a number of criteria, including the beneficial agent potency and concentration, and the process by which the formulation is prepared. The viscosity of the formulation can be chosen so that the desired amount of beneficial agent is delivered over the desired period of time.

The invention also consists of non-aqueous single phase biocompatible viscous vehicles capable of uniformly suspending beneficial agents and formulations containing at least one beneficial agent uniformly suspended in said viscous vehicle. The invention also consists of formulations containing at least one beneficial agent uniformly suspended in a non-aqueous single phase biocompatible viscous vehicle, which formulations are stable for an extended period of time at body temperatures, and capable of delivering said beneficial agents uniformly at low flow rates. The discovery consists of the realization that stable non-aqueous viscous vehicles improve the stability of beneficial agents in a wide range of formulation conditions including concentration, elevated temperatures and duration of stable formulation, thus making possible the delivery of beneficial agents in long term implantable devices that would not otherwise be feasible.

#### Definitions

As used herein, the following terms have the following meanings:

The term "chemical stability" means that an acceptable percentage of degradation products produced by chemical pathways such as oxidation, deamidation, or hydrolysis is formed. In particular, a formulation is considered chemically stable if no more than about 35% breakdown products are formed after 2 months at 37°C.

The term "physical stability" means that an acceptable percentage of aggregates (e.g., dimers, trimers and larger forms) are formed by the beneficial agent. For the formulation (viscous vehicle and beneficial agent) this term means that the formulation retains stability, flowability, and the ability to uniformly dispense the beneficial agent. In particular, a formulation is considered physically stable if no more than about 15% aggregates are formed after two months at 37°C.

The term "stable formulation" means that at least about 65% chemically and physically stable beneficial agent remains after two months at 37°C (or equivalent conditions at an elevated temperature). Particularly preferred formulations are those which retain at least about 80% chemically and physically stable beneficial agent under these conditions. Especially preferred stable formulations are those which do not exhibit degradation after sterilizing irradiation (e.g., gamma, beta or electron beam).

The term "beneficial agent" means peptides, proteins, nucleotides, hormones, viruses, antibodies, etc. that comprise polymers of amino acid or nucleic acid residues. These beneficial agents are generally degradable in water and generally stable as a dry powder at elevated temperatures. Synthetically produced, naturally derived or recombinantly produced moieties are included in this term. The term also includes lipoproteins and post translationally modified forms, e.g., glycosylated proteins. Analogs, derivatives, agonists, antagonists and pharmaceutically acceptable salts of any of these are included in this term. The term also includes proteins and/or protein substances which have D-amino acids, modified, derivatized or non-naturally occurring amino acids in the D- or L- configuration and/or peptomimetic units as part of their structure. The term protein will be used in

the present invention. The term also means that the beneficial agent is present in the solid state, e.g., powder or crystalline.

The term "excipient" means a more or less inert substance in a formulation that is added as a diluent or vehicle or to give form or consistency.

5 Excipients are distinguished from solvents such as ETOH, which are used to dissolve drugs in formulations. Excipients include non-ionic surfactants such as polysorbates, which are used to solubilize drugs in formulations; preservatives such as benzyl alcohols or methyl or propyl parabens, which are used to prevent or inhibit microbial growth; chelating agents; flavoring  
10 agents; and other pharmaceutically acceptable formulation aides.

The term "viscous vehicle" means a vehicle with a viscosity in the range of about 1,000 to 10,000,000 poise. The term includes Newtonian and non-Newtonian materials. Preferred are vehicles with a viscosity of about 10,000 to 250,000 poise. The formulations of this invention can uniformly  
15 expel beneficial agents suspended in the viscous vehicle from implantable drug delivery devices. The formulations exhibit a shear rate at the exit of said devices of 1 to  $1 \times 10^{-7}$  reciprocal second, preferably an exit shear rate of  $1 \times 10^{-2}$  to  $1 \times 10^{-5}$  reciprocal second.

The term "single phase" means a solid, semi-solid, or liquid  
20 homogeneous system that is both physically and chemically uniform throughout as determined by differential scanning calorimetry (DSC). The DSC scan should show one peak indicative of a single phase.

The term "biocompatible" means a property or characteristic of a viscous vehicle to disintegrate or break down, over a prolonged period of time,  
25 in response to the biological environment in the patient, by one or more physical or chemical degradative processes, for example by enzymatic action, oxidation or reduction, hydrolysis (proteolysis), displacement, e.g. ion exchange, or dissolution by solubilization, emulsion or micelle formation, and which material is then absorbed by the body and surrounding tissue, or  
30 otherwise dissipated thereby.

The term "polymer" includes polyesters such as PLA (polylactic acid) [having an inherent viscosity in the range of about 0.5 to 2.0 i.v.] and PLGA

(polylacticpolyglycolic acid) [having an inherent viscosity in the range of about 0.5 to 2.0 i.v.], pyrrolidones such as polyvinylpyrrolidone (having a molecular weight range of about 2,000 to 1,000,000), esters or ethers of unsaturated alcohols such as vinyl acetate, and polyoxyethylenepolyoxypropylene block copolymers (exhibiting a high viscosity at 37°C) such as Pluronic 105. Currently preferred polymer is polyvinylpyrrolidone.

The term "solvent" includes carboxylic acid esters such as lauryl lactate, polyhydric alcohols such as glycerin, polymers of polyhydric alcohols such as polyethylene glycol (having a molecular weight of about 200 to 600), fatty acids such as oleic acid and octanoic acid, oils such as castor oil, propylene carbonate, lauryl alcohol, or esters of polyhydric alcohols such as triacetin acetate. Currently preferred is lauryl lactate.

The term "surfactant" includes esters of polyhydric alcohols such as glycerol monolaurate, ethoxylated castor oil, polysorbates, esters or ethers of saturated alcohols such as myristyl lactate (Ceraphyl 50), and polyoxyethylenepolyoxypropylene block copolymers such as Pluronic. Currently preferred are glycerol monolaurate and polysorbates.

The term "antioxidant" means a pharmaceutically acceptable aid for stabilization of the beneficial agent against degradation such as oxidation. Antioxidants include, but are not limited to, tocopherol (vitamin E), ascorbic acid, ascorbyl palmitate, butylated hydroxyanisole, butylated hydroxytoluene, and propyl gallate. A preferred antioxidant depends on solubility and the efficiency of the antioxidant for protecting against degradation or chemical change of the beneficial agent in the preferred vehicle. Currently preferred is ascorbyl palmitate.

#### Preparation of Formulations

The present invention is drawn to stable non-aqueous single phase biocompatible viscous vehicles capable of suspending beneficial agents and uniformly dispensing said beneficial agents at body temperatures at low flow rates over an extended period of time. The present invention is also directed to formulations containing beneficial agents uniformly suspended in said

single phase biocompatible viscous vehicles which are stable for prolonged periods of time at body temperatures.

Examples of beneficial agents that may be formulated using the present invention include those peptides or proteins that have biological activity or that may be used to treat a disease or other pathological condition. They include, but are not limited to, adrenocorticotrophic hormone, angiotensin I and II, atrial natriuretic peptide, bombesin, bradykinin, calcitonin, cerebellin, dynorphin N, alpha and beta endorphin, endothelin, enkephalin, epidermal growth factor, fertirelin, follicular gonadotropin releasing peptide, galanin, glucagon, GLP-1, gonadorelin, gonadotropin, goserelin, growth hormone releasing peptide, histrelin, human growth hormone, insulin, interferons, leuprolide, LHRH, motilin, nafarerlin, neurotensin, oxytocin, relaxin, somatostatin, substance P, tumor necrosis factor, triptorelin, vasopressin, growth hormone, nerve growth factor, blood clotting factors, ribozymes, and antisense oligonucleotides. Analogs, derivatives, antagonists agonists and pharmaceutically acceptable salts of the above may also be used.

The beneficial agents useful in the formulations and methods of the present invention can be used in the form of a salt, preferably a pharmaceutically acceptable salt. Useful salts are known to those of skill in the art and include salts with inorganic acids, organic acids, inorganic bases, or organic bases.

Beneficial agents that are not readily soluble in non-aqueous solvents are preferred for use in the present invention. One of skill in the art can easily determine which compounds will be useful on the basis of their solubility. The amount of beneficial agent may vary depending on the potency of the compound, the condition to be treated, the solubility of the compound, the expected dose and the duration of administration. (See, for example, Gilman, et. al, The Pharmacological Basis of Therapeutics, 7<sup>th</sup> ed. (1990) and Remington, Pharmacological Sciences, 18<sup>th</sup> ed. (1990), the disclosures of which are incorporated herein by reference.)

It has been unexpectedly found that using a stable non-aqueous single phase biocompatible viscous vehicle increases the stability of the beneficial



agent. For example, as seen in Figures 1 and 2, human growth hormone (hGH) was found to be stable at 37°C over 12 weeks in formulations of polyvinylpyrrolidone/PEG; Pluronic; and glycerol monolaurate/lauryl lactate/polyvinylpyrrolidone. Figure 1 shows stability results using reverse phase  
5 HPLC. Figure 2 shows stability results using size exclusion chromatography.

Generally, stable non-aqueous single phase biocompatible viscous vehicles may be prepared by combining the dry (low moisture content) ingredients in a dry box or under other dry conditions and blending them at elevated temperature, preferably about 40 to about 70°C, to allow them to  
10 liquify. The liquid vehicle is allowed to cool to room temperature. Differential scanning calorimetry was used to verify that the vehicle was single phase. The final moisture content of the viscous vehicle was <2%.

Generally, the stable formulations of the present invention may be prepared by combining the vehicle and beneficial agent under dry conditions and blending them under vacuum at elevated temperature, preferably about  
15 40 to about 70°C, to disperse the beneficial agent uniformly throughout the vehicle. The formulation is allowed to cool to room temperature.

It has been found that drying the beneficial agent prior to formulation enhances the stability of the formulation.

20 It has also been found that adding antioxidants, such as tocopherol, ascorbic acid, ascorbyl palmitate, butylated hydroxyanisole, butylated hydroxytoluene, and propyl gallate reduces the formation of degradation products (e.g., unstable chemical intermediates) during sterilization.

## 25 Methodology

We have found that stable non-aqueous beneficial agent formulations utilizing viscous vehicles may be prepared by combining the ingredients for the viscous vehicle under dry conditions and blending them at elevated temperature to allow them to liquify and form a single phase. Once a single  
30 phase viscous vehicle is formed, the vehicle is allowed to cool to room temperature. Beneficial agent is added with mixing at elevated temperature under vacuum to uniformly disperse it in the viscous vehicle.

We have tested these beneficial agent formulations, for example formulations of hGH, for stability by subjecting them to accelerated aging tests. Results show that these formulations remained stable over extended periods of time.

5 We have tested beneficial agent formulations, for example human growth hormone and lysozyme, for stability by suspending them in a variety of non-aqueous single phase viscous vehicles prepared according to the present invention, then subjecting them to accelerated aging at elevated temperatures. The stability of the formulations was measured. Results of  
10 these studies demonstrate that these formulations were stable at conditions that approximate or exceed storage for one year at 37°C.

We have also tested beneficial agent formulations prepared as described herein for stability after 2.5 megarads gamma irradiation. Results show that these formulations remained chemically and physically stable after  
15 such irradiation.

#### Methods

The following methods were used to perform the studies in the Examples that follow.

##### 20 1. Preparing Protein powders

Human Growth Hormone (obtained for example, from BresaGen Limited, Adelaide, Australia)

The active agent was reconstituted in deionized water. The solution containing the active agent was buffer exchanged using an Amicon Diaflo®  
25 Ultrafiltration membrane (molecular weight cut-off 10,000).

The diafiltrated active agent solution was spray dried using a Yamato mini-spray dryer. Powder was collected in a collection vessel through a cyclone trap. All handling of the spray dried powder took place in a dry box evacuated with nitrogen. The generated powder was analyzed for particle  
30 size and distribution, moisture content, protein content and stability by size exclusion and reverse-phase chromatography.

It is known that the conformation of some proteins can be stabilized by the addition of a sugar (such as sucrose or mannitol) or a polyol (such as ethylene glycol, glycerol, glucose, and dextran.)

## 2. Preparation of Viscous Vehicles

5 We have found that stable single phase biocompatible viscous vehicles may be prepared by combining the ingredients and blending them at elevated temperatures to allow them to liquify and form a single phase. A differential scanning calorimetry scan showed one peak, indicative of a single phase. The mixing was completed under vacuum to remove trapped air bubbles  
10 produced from the powders. The mixer was a dual helix blade mixer (D.I.T.) which runs at a speed around 40 rpm. Higher speeds can be used but are not required.

If a three component viscous vehicle is prepared, the solvent portion of the vehicle was added to the heated bowl of the mixer first, followed by the  
15 surfactant. The polymer was added last, and the ingredients were mixed until a solution (single phase) resulted. Vacuum was applied during mixing to remove air bubbles. The solution was dispensed from the bowl while at elevated temperature, and allowed to cool to room temperature. On cooling the vehicle exhibited increased viscosity. Two and single component gels  
20 were made using the same process.

## 3. Preparation of beneficial agent formulations

To prepare the formulation, the single phase viscous vehicle was heated and then blended under vacuum with a weighed amount of beneficial  
25 agent. The beneficial agent and the single phase viscous vehicle were blended in the same manner as the vehicle was prepared, using a dual helix blade mixer (or other similar mixer). Mixing speed was between 40 and 120 rpm for approximately 15 minutes or until a uniform dispersion was attained. The resulting mixture was removed from the mixer, sealed in a dry container,  
30 and allowed to cool to room temperature.

4. Preparation of reservoirs

The reservoirs of implantable drug delivery devices (as disclosed in U.S. Patent Application Serial No. 08/595,761, incorporated herein by reference) were filled with the appropriate hGH formulation. The formulation was filled into titanium reservoirs with a polymer plug blocking each end. The filled reservoir was then sealed in a polyfoil bag and placed in a stability testing oven.

It should be noted that the formulations in the reservoirs of these devices are completely isolated from the outside environment.

5. Reverse Phase-HPLC (RP-HPLC)

All stability samples of hGH were assayed for protein content and chemical stability by reverse phase chromatography (RP-HPLC). Analyses were performed on a Hewlett Packard HP-1090 system with a refrigerated autosampler (4°C). The chromatographic conditions used are listed below.

TABLE 1  
RP-HPLC Chromatographic Conditions

Description	Parameter		
Column	J.T. Baker-C18, 4.6x250 mm		
Flow Rate	1.0 mL/min		
Detection	214 nm		
Mobile Phase	A: 0.1% TFA in water B: 0.1% TFA in acetonitrile		
Gradient	<u>time</u>	<u>%A</u>	<u>%B</u>
	0	65	35
	5	50	50
	45	35	65
	50	30	70
	55	65	35

An hGH reference standard solution was prepared and its protein content calculated from the absorbance measurement at 280 nm. Three dilutions of this solution, representing 80%, 100%, and 120% of the expected concentration of hGH in the samples were run in duplicate at the beginning and the end of each run and used to calculate total protein content of the samples.

#### 6. Size Exclusion Chromatography (SEC)

All stability samples of hGH were assayed for protein content and high molecular weight degradation products by size exclusion chromatography. Analyses were performed on a Hewlett Packard HP-1090 system with a refrigerated autosampler (4°C). The chromatographic conditions used are listed below

TABLE 2  
SEC Chromatographic Conditions

Description	Parameter
Column	TSK-2000SWXL
Flow Rate	0.5 ml/ min
Detection	214 nm
Mobile Phase	25 mM sodium phosphate, 100 mM sodium chloride, pH 7.0

A hGH reference standard solution was prepared and its protein content calculated from the absorbance measurement at 280 nm. Three dilutions of this solution, representing 80%, 100%, and 120% of the expected concentration of hGH in the samples were run in duplicate at the beginning and the end of each run and used to calculate total protein content of the samples. The amount of high molecular weight degradation products was calculated by area normalization.

The following examples are offered to illustrate this invention and are not meant to be construed in any way as limiting the scope of this invention.

EXAMPLE 1Preparation of Non-aqueous Single Phase Viscous Vehicles

The non-aqueous single phase viscous vehicles can be prepared as follows and shown in the below table

- 5
- A. Glycerol monolaurate (Danisco Ingredients, New Century, Kansas) (25 g) was dissolved in lauryl lactate (ISP Van Dyk Inc., Belleville, NJ) (35 g) at 65°C. Polyvinylpyrrolidone C30 (BASF, Mount Olive, NJ) (40 g) was added and the mixture blended at about 40 rpm in a dual helix blade mixer (D.I.T.) until a single phase was achieved. Trapped air bubbles were removed by applying vacuum to the mixing chamber. The single phase vehicle was dispensed from the mixer, and allowed to cool to room temperature.
- 10
- B. Glycerol monolaurate (Danisco Ingredients, New Century, Kansas) (25 g) was dissolved in lauryl lactate (ISP Van Dyk Inc., Belleville, NJ) (35 g) at 65°C. Polyvinylpyrrolidone C17 (BASF, Mount Olive, NJ) (40 g) was added and the mixture blended at about 40 rpm in a dual helix blade mixer (D.I.T.) until a single phase was achieved. Trapped air bubbles were removed by applying vacuum to the mixing chamber. The single phase vehicle was dispensed from the mixer, and allowed to cool to room temperature.
- 15
- 20
- C. Polyvinylpyrrolidone C30 (BASF, Mount Olive, NJ) (50 g) was dissolved in polyethylene glycol 400 (Union Carbide) (50 g) at approximately 65°C until a single phase solution was formed. The single phase vehicle was dispensed from the mixer, and allowed to cool to room temperature.
- 25
- D. Polyvinylpyrrolidone C17 (BASF, Mount Olive, NJ) (50 g) was dissolved in polyethylene glycol 400 (Union Carbide) (50 g) at approximately 65°C until a single phase solution was formed. The single phase vehicle was dispensed from the mixer, and allowed to cool to room temperature.
- 30

- 5 E. Polyvinylpyrrolidone C17 (BASF, Mount Olive, NJ) (50 g) was dissolved in castor oil (Spectrum, Gardena, CA) (50 g) at approximately 65°C until a single phase solution was formed. The single phase vehicle was dispensed from the mixer, and allowed to cool to room temperature.
- 10 F. Polyvinylpyrrolidone C17 (BASF, Mount Olive, NJ) (50 g) was dissolved in octanoic acid (Spectrum, Gardena, CA) at approximately 65°C until a single phase solution was formed. The single phase vehicle was dispensed from the mixer, and allowed to cool to room temperature.
- G. Polyvinylpyrrolidone C17 (BASF, Mount Olive, NJ) (50 g) was dissolved in oleic acid (Spectrum, Gardena, CA) at approximately 65°C until a single phase solution was formed. The single phase vehicle was dispensed from the mixer, and allowed to cool to room temperature.
- 15 H. Polyvinylpyrrolidone C17 (BASF, Mount Olive, NJ) (35%) was dissolved in glycerin (Baker, NJ) (65%) at approximately 65°C until a single phase solution was formed. The single phase vehicle was dispensed from the mixer, and allowed to cool to room temperature.
- 20 I. Cremophor EL (ethoxylated castor oil) (BASF, Mount Olive, NJ) (5%) was dissolved in castor oil (Spectrum, Gardena, CA) (70%), and polyvinylpyrrolidone C17 (BASF, Mount Olive, NJ) (25%) was added and dissolved by mixing at approximately 40 rpm to form a single phase vehicle. The single phase vehicle was dispensed from the mixer, and allowed to cool to room temperature.
- 25 J. Pluronic 105 (BASF, Mount Olive, NJ) was heated to approximately 65°C with mixing until melted. The single phase vehicle was dispensed from the mixer, and allowed to cool to room temperature.

Table 3  
Component Ratios

	Polymer	Component		Ratio	Viscosity at Low
		Surfactant	Solvent		Shear Rate (Poise)
5	PVP	GML	LL	53:5:42	25,000
	PVP	GML	LL	55:10:35	50,000
	PVP	GML	LL	50:15:35	7,000
	PVP	-----	LA	60:40	
	PVP	Ceraphyl 50	LA	60:10:30	
10	PVP	-----	oleic acid	50:50	30,000
	PVP	-----	octanoic acid	55:45	7,000
	PVP	polysorbate 80	-----	50:50	
	PVP	-----	PEG 400	50:50	
	PVP	caster oil	-----	50:50	
15	-----	Pluronic 105	-----	100	1,000,000
	PVP	-----	glycerin	50:50	5,000

Wherein:

GML = glycerol monolaurate

20 LL = lauryl lactate

PVP = polyvinylpyrrolidone C30

LA = lauryl alcohol

PEG = polyethyleneglycol 400

25

## EXAMPLE 2

### Preparation of hGH

#### A. Preparation by spray drying

Lyophilized hGH (BresaGen Limited, Adelaide, Australia) was  
30 reconstituted in 150 ml of deionized water. This stock solution contained  
1050 mg of hGH. Buffer exchange was accomplished using an Amicon  
Diaflo® Ultrafiltration membrane (molecular weight cut-off 10,000). The



ultrafiltration cell was connected to an auxilliary reservoir containing 5mM phosphate buffer (pH 7). The cell's fluid volume, as well as the hGH concentration, remained constant as excipients were replaced by phosphate buffer.

5           The diafiltrated protein solution (protein concentration in the solution approximately 2%) was spray dried using a Yamato mini-spray dryer. Settings on the spray dryer were as follows: aspiration pressure constantly adjusted to 1.3 kgf/cm<sup>2</sup>, inlet temperature 120°C, solution flow rate 2.5 (approximately 3 ml/min). Powder was collected in a collection vessel through  
10 a cyclone trap. All handling of the spray dried powder took place in a dry box evacuated with nitrogen (% RH: 1-4%). The water content of the suspending vehicles is shown in the below table.

TABLE 4  
WATER CONTENT OF SUSPENDING VEHICLES

Vehicle	Water Content of Vehicle at T 0 % w/w	Water Content of Vehicle in 12 wks. At 37°C % w/w
Pluronic 105	0.25	0.4
GML/LL/PVP	1.5	1.3
PVP/PEG	2.0	2.0

25   Wherein:

GML = glycerol monolaurate

LL = lauryl lactate

PVP = polyvinylpyrrolidone C30

PEG = polyethyleneglycol 400

30

EXAMPLE 3  
Preparation of hGH Formulation

A portion of the single phase viscous vehicle was weighed (9 g) and  
5 heated to 60°C. hGH (BresaGen Limited, Adelaide, Australia) (1 g) was  
added to the vehicle and mixed for 15 minutes. The mixing was completed  
under vacuum to remove air bubbles added from the powder.

Approximately 10 mg of the spray-dried hGH powder were weighed out  
(content of hGH in the powder was recalculated based on the determined  
10 water and salt content) and mixed with 100 µl of the vehicle at 55-65°C (3  
samples per each vehicle). Special care was taken while mixing powder in  
the suspending vehicle to achieve maximum particle uniform dispersion in the  
vehicle. All steps were done in a dry box.

The resulting suspension was dissolved with 10 ml of release rate  
15 buffer and analyzed by size exclusion and reverse-phase chromatography.  
Spray dried hGH powder was used as a control.

TABLE 5  
STABILITY OF hGH SUSPENSIONS AT 37°C AS MEASURED BY  
20 SIZE EXCLUSION CHROMATOGRAPHY

	Time Weeks	Spray-dried Powder -80°C	PVP/PEG 400 suspension	GML/LL/PVP suspension	Pluronic 105 suspension
		%LS	%LS	%LS	%LS
25	0	96±1	88±6	92±2	87±7
	1	99±8	81±2	94±3	93±3
	2	99±3	83±1	97±1	94±1
	3	97±1	84±2	95±2	95±3
30	4	95±2	82±8	94±4	93±5
	7	95±4	76±3	93±4	88±2
	12	97±4	79±3	97±1	95±6

Each data point represents the mean ± relative standard deviation of three individual samples taken from three separate vials.

TABLE 6  
STABILITY OF hGH SUSPENSIONS at 37°C AS MEASURED BY  
REVERSE PHASE CHROMATOGRAPHY

Time Weeks	spray-dried Powder -80°C %LS	PVP/PEG 400 suspension %LS	GML/LL/PVP suspension %LS	Pluronic 105 suspension %LS
0	104±1	99±3	99±2	89±7
1	104±8	78±2	98±3	96±6
2	104±4	73±3	95±1	96±1
3	104±2	78±4	97±3	97±4
4	100±2	74±10	93±4	96±4
7	108±5	72±4	96±2	94±2
9	102±3	66±3	92±3	93±2
12	101±2	66±1	89±2	92±5

Each data point represents the mean ± relative standard deviation of three individual samples taken from three separate vials.

EXAMPLE 4  
Preparation of Reservoirs  
Release Rate Profiles

Titanium reservoir systems of implantable drug delivery devices (as disclosed in U.S. Patent Application Serial No. 08/595,761, incorporated herein by reference) were each assembled with an osmotic engine, piston, and rate controlling membrane. The reservoirs were filled with the appropriate amount of viscous vehicle formulation and capped with a flow plug. The systems were placed in a water bath at 37°C, and allowed to

release formulation for an extended period of time. Released material was sampled twice per week. Assays for released material were completed using reverse phase HPLC. The resulting concentrations of beneficial agent for each system were converted to released amount per day. The beneficial agent was found to have a zero order release from the implantable drug delivery device. As shown in Figures 3 through 8.

EXAMPLE 5

Stability of hGH in Non-aqueous Viscous Vehicle Formulations

Formulations of 10% w/w hGH in vehicle were prepared as described above and placed in vials. The formulations were subjected to accelerated aging by storing them at elevated temperatures and times shown in the below table in a controlled temperature oven .

TABLE 7

Vehicle	Time(hrs)	Temperature	%LS by SEC	%LS by RP-HPLC
Pluronic 105	0	50°C	98±3	101±3
Pluronic 105	1	50°C	98±3	101±4
Pluronic 105	2	50°C	100±1	102±3
Pluronic 105	4	50°C	101±3	105±3
GML/LL/PVP	0	65°C	99±3	101±3
GML/LL/PVP	1	65°C	93±6	97±6
GML/LL/PVP	2	65°C	91±5	95±5
GML/LL/PVP	4	65°C	95±3	98±3

Each data point represents the mean ± relative standard deviation of three individual samples taken from three separate vials.

Results, presented in the following table, demonstrate that these formulations were able to maintain the stability of the hGH in each case. In each case, at least 70% hGH was retained.

5

TABLE 8

## RECOVERY OF hGH FROM NONAQUEOUS SUSPENSIONS

	<u>Vehicle</u>	<u>%LS by RP-HPLC</u>	<u>%LS by Size-exclusion HPLC</u>
	PVP/PEG 400	99±3%	88±6%
10	GML/LL/PVP	99±2%	92±2%
	Pluronic 105	89±7%	87±7%

Each data point represents the mean  $\pm$  relative standard deviation of three individual samples taken from three separate vials.

15 %LS or % label strength = (measured protein content + theoretical protein content) x 100%

Modification of the above-described modes of carrying out various embodiments of this invention will be apparent to those of skill in the art following the teachings of this invention as set forth herein. The examples  
20 described above are not limiting, but are merely exemplary of this invention, the scope of which is defined by the following claims.

## Claims

1. A stable non-aqueous single phase biocompatible viscous vehicle capable of suspending beneficial agents and homogeneously dispensing said  
5 beneficial agent over an extended period of time at body temperature and at low flow rates.
2. The vehicle of claim 1 comprising two components selected from the group consisting of solvent, surfactant, and polymer, wherein the two  
10 components are not of the same type.
3. The vehicle of claim 1 comprising at least two components selected from the group consisting of solvent, surfactant, and polymer, wherein the components are not of the same type.  
15
4. The vehicle of claim 1 which comprises three components selected from the group consisting of solvent, surfactant, and polymer, wherein the components are not of the same type.
- 20 5. The vehicle of claim 2 or 4 wherein said solvent is selected from the group carboxylic acid esters, polyhydric alcohols, polymers of polyhydric alcohols, fatty acids, oils, propylene carbonate, lauryl alcohol, and esters of polyhydric alcohols.
- 25 6. The vehicle of claim 2 or 4 wherein said surfactant is selected from the group esters of polyhydric alcohols, ethoxylated castor oil, polysorbates, esters or ethers of saturated alcohols, and polyoxyethylenepolyoxypropylene block copolymers.
- 30 7. The vehicle of claim 2 or 4 wherein said polymer is selected from the group polyesters, pyrrolidones, esters or ethers of unsaturated alcohols, and polyoxyethylenepolyoxypropylene block copolymers.

8. The vehicle of claim 2 wherein the ratios of the components are in the range of 40:60 to 60:40.

5 9. The vehicle of claim 4 wherein the ratios of the components are in the range of about 30% to about 50% for solvent, about 5% to about 20% for surfactant, and about 5% to about 60% for polymer.

10 10. The vehicle of claim 4 wherein the polymer is polyvinylpyrrolidone, the surfactant is gml, and the solvent is lauryl lactate.

11. The vehicle of claim 4 wherein the polymer is polyvinylpyrrolidone, the surfactant is polysorbate, and the solvent is lauryl lactate.

15 12. The vehicle of claim 1 which comprises an antioxidant.

13. The vehicle of claim 12 wherein said antioxidant is selected from the group consisting of tocopherol, ascorbic acid, ascorbyl palmitate, butylated hydroxyanisole, butylated hydroxytoluene, and propyl gallate.

20 14. A stable non-aqueous viscous protein formulation comprising  
a) at least one beneficial agent, and  
b) a non-aqueous single phase biocompatible viscous vehicle, which formulation is capable of being uniformly dispensed over an extended period of time at a low flow rate.

25 15. A non-aqueous formulation comprising at least one beneficial agent uniformly suspended in a non-aqueous single phase biocompatible viscous vehicle, which formulation can be delivered from an implantable drug delivery system such that the exit shear rate of the formulation is between about 1 and  
30  $1 \times 10^{-7}$  reciprocal second.

16. The formulation of claim 14 wherein said formulation is stable at body temperature for extended periods of time.

17. The formulation of claim 14 which comprises at least about 0.1% (w/w) beneficial agent.
- 5 18. The formulation of claim 14 which comprises at least about 10% (w/w) beneficial agent.
19. The formulation of claim 14 wherein said beneficial agent is selected from the group consisting of peptide, protein, nucleotide, hormone, virus, or  
10 antibody.
20. The formulation of claim 19 wherein said beneficial agent is a protein.
21. The formulation of claim 14 which is stable at 65° C for at least about 2  
15 months.
22. The formulation of claim 14 which is stable at 37° C for at least about 3 months.
- 20 23. The formulation of claim 14 which is stable at 37° C for at least about one year.
24. The formulation of claim 14 which is adapted for use in an implantable drug delivery device.  
25
25. The formulation of claim 14 wherein said vehicle is selected from the group consisting of solvent, surfactant and polymer.
26. The formulation of claim 14 wherein said vehicle comprises an  
30 antioxidant.
27. The formulation of claim 14 comprising a beneficial agent which has been dried to a low moisture content prior to incorporation in said formulation.



28. The formulation of claim 14 which is stable after sterilization.
29. A method for preparing the stable single phase viscous vehicle of claim  
5 1 comprising the steps of (1) blending the ingredients at elevated temperature under dry conditions to allow them to liquify, and (2) allowing the liquid from step (1) to cool to room temperature.
30. A method for preparing the stable formulation of claim 14 comprising  
10 combining the single phase viscous vehicle and beneficial agent under dry conditions and blending them under vacuum at elevated temperature to uniformly disperse the beneficial agent in the vehicle, and allowing the formulation to cool to room temperature.
31. The method of claim 30 wherein at least about 0.1% (w/w) beneficial  
15 agent is suspended in said vehicle.
32. The method of claim 30 wherein at least about 10% (w/w) beneficial  
20 agent is suspended in said vehicle.
33. A method for treating a subject suffering from a condition which may be alleviated by administration of a beneficial agent comprising administering to said subject a therapeutically effective amount of the formulation of Claim  
25 14.
34. The method of claim 33 wherein said administration is parenteral administration.
35. The method of claim 33 wherein said administration is long-term  
30 continuous administration.
36. The method of claim 33 wherein said administration is accomplished by use of an implantable drug delivery system.

37. The method of claim 33 wherein said daily administration continues for a period selected from the group consisting of about 3 months, about 6 months, and about 12 months.

5

38. The method of claim 37 wherein said daily administration is accomplished using an implantable drug delivery system.

Approved for release

## 5

10

[illegible]




FIGURE 1






FIGURE 2




## FIGURE 3

**IN VITRO SYSTEMS, PRE-RAT STUDY**  
**AVERAGE RELEASE RATE ( $\mu\text{g/day}$ )**



**FIGURE 4**



## FIGURE 5







**45% LOADED PARTICLES IN GML:LL:PVP VEHICLE**  
**RELEASE RATE ( $\mu\text{g/day}$ ), n=4**

**RELEASE RATE ( $\mu\text{g/day}$ )**

**TIME (days)**

**TARGET: 819  $\mu\text{g/day}$**

TIME (days)	RELEASE RATE ( $\mu\text{g/day}$ )
0	1000
2	1000
4	1050
6	950
8	900
10	880
12	860
14	850
16	840
18	830
20	820
22	810

Docket No.  
2914R1

# Declaration and Power of Attorney For Patent Application

## English Language Declaration

As a below named inventor, I hereby declare that:

My residence, post office address and citizenship are as stated below next to my name,

I believe I am the original, first and sole inventor (if only one name is listed below) or an original, first and joint inventor (if plural names are listed below) of the subject matter which is claimed and for which a patent is sought on the invention entitled

**STABLE NON-AQUEOUS SINGLE PHASE VISCOUS VEHICLES AND FORMULATIONS UTILIZING SUCH VEHICLES**

the specification of which

(check one)

☒ is attached hereto.

☐ was filed on \_\_\_\_\_ as United States Application No. or PCT International Application Number \_\_\_\_\_ and was amended on \_\_\_\_\_ (if applicable)

I hereby state that I have reviewed and understand the contents of the above identified specification, including the claims, as amended by any amendment referred to above.

I acknowledge the duty to disclose to the United States Patent and Trademark Office all information known to me to be material to patentability as defined in Title 37, Code of Federal Regulations, Section 1.56.

I hereby claim foreign priority benefits under Title 35, United States Code, Section 119(a)-(d) or Section 365(b) of any foreign application(s) for patent or inventor's certificate, or Section 365(a) of any PCT International application which designated at least one country other than the United States, listed below and have also identified below, by checking the box, any foreign application for patent or inventor's certificate or PCT International application having a filing date before that of the application on which priority is claimed.

Prior Foreign Application(s)

Priority Not Claimed

\_\_\_\_\_  
(Number)

\_\_\_\_\_  
(Country)

\_\_\_\_\_  
(Day/Month/Year Filed)

☐

\_\_\_\_\_  
(Number)

\_\_\_\_\_  
(Country)

\_\_\_\_\_  
(Day/Month/Year Filed)

☐

\_\_\_\_\_  
(Number)

\_\_\_\_\_  
(Country)

\_\_\_\_\_  
(Day/Month/Year Filed)

☐

I hereby claim the benefit under 35 U.S.C. Section 119(e) of any United States provisional application(s) listed below:

<u>60/119,170</u>	<u>2/8/99</u>
(Application Serial No.)	(Filing Date)
(Application Serial No.)	(Filing Date)
(Application Serial No.)	(Filing Date)

I hereby claim the benefit under 35 U. S. C. Section 120 of any United States application(s), or Section 365(c) of any PCT International application designating the United States, listed below and, insofar as the subject matter of each of the claims of this application is not disclosed in the prior United States or PCT International application in the manner provided by the first paragraph of 35 U.S.C. Section 112, I acknowledge the duty to disclose to the United States Patent and Trademark Office all information known to me to be material to patentability as defined in Title 37, C. F. R., Section 1.56 which became available between the filing date of the prior application and the national or PCT International filing date of this application:

(Application Serial No.)	(Filing Date)	(Status) (patented, pending, abandoned)
(Application Serial No.)	(Filing Date)	(Status) (patented, pending, abandoned)
(Application Serial No.)	(Filing Date)	(Status) (patented, pending, abandoned)

I hereby declare that all statements made herein of my own knowledge are true and that all statements made on information and belief are believed to be true; and further that these statements were made with the knowledge that willful false statements and the like so made are punishable by fine or imprisonment, or both, under Section 1001 of Title 18 of the United States Code and that such willful false statements may jeopardize the validity of the application or any patent issued thereon.

POWER OF ATTORNEY: As a named inventor, I hereby appoint the following attorney(s) and/or agent(s) to prosecute this application and transact all business in the Patent and Trademark Office connected therewith. *(list name and registration number)*


Steven F. Stone	Reg. No. 20,246
Paul L. Sabatine	Reg. No. 22,539
D. Byron Miller	Reg. No. 30,661
John A. Dhuey	Reg. No. 26,265
Pauline A. Clarke	Reg. No. 29,783
Owen Bates	Reg. No. 40,436
Susan K. Thomas	Reg. No. 39,780


Send Correspondence to:

**Pauline A. Clarke**  
**ALZA Corporation, 1900 Charleston Road, P.O. Box 7210**  
**Mt. View, CA 94039-7210**

Direct Telephone Calls to: *(name and telephone number)*

**Pauline A. Clarke 650-564-5560**

Full name of sole or first inventor <b>Stephen A. Berry</b>	
Sole or first inventor's signature 	Date <b>1/25/00</b>
Residence <b>1050 Spring Grove Road, Hollister, CA 95023</b>	
Citizenship <b>USA</b>	
Post Office Address <b>Same</b>	

Full name of second inventor, if any <b>Pamela J. Ferreira</b>	
Second inventor's signature 	Date <b>1/25/00</b>
Residence <b>304 CanoeCourt, Redwood City, CA 94065</b>	
Citizenship <b>USA</b>	
Post Office Address <b>Same</b>	

Full name of third inventor, if any <b>Houdin Dehnad</b>	
Third inventor's signature <i>Houdin Dehnad</i>	Date 01/25/2000
Residence <b>507 Vallejo Street, El Granada, CA 94018</b>	
Citizenship <b>Iran</b>	
Post Office Address <b>Same</b>	

Full name of fourth inventor, if any <b>Anna Muchnik</b>	
Fourth inventor's signature <i>Anna Muchnik</i>	Date 01/26/00
Residence <b>1542 Winding Way, Belmont, CA 94002</b>	
Citizenship <b>USA</b>	
Post Office Address <b>Same</b>	

Full name of fifth inventor, if any	
Fifth inventor's signature	Date
Residence	
Citizenship	
Post Office Address	

Full name of sixth inventor, if any	
Sixth inventor's signature	Date
Residence	
Citizenship	
Post Office Address	